Association for Information Systems AIS Electronic Library (AISeL)

BLED 2014 Proceedings

BLED Proceedings

6-2014

Microlearning mApp to Improve Long Term Health Behaviours: Design and Test of Multi-Channel Service Mix

Luuk P.A. Simons

Delft University of Technology, Netherlands, L.P.A.Simons@tudelft.nl

Florian Foerster

Georgia Institute of Technology, USA, florian.foerster@gatech.edu

Peter A. Bruck

Research Studio Micro-learning and Information Environment; Research Studios Austria Forschungsgesellschaft, Austria, bruck@researchstudio.at

Luvai Motiwalla

Operations & Info Systems Dept, U Massachusetts Lowell, USA, luvai_motiwalla@uml.edu

Catholijn M. Jonker

Delft University of Technology, Netherlands, c.m.jonker@tudelft.nl

Follow this and additional works at: http://aisel.aisnet.org/bled2014

Recommended Citation

Simons, Luuk P.A.; Foerster, Florian; Bruck, Peter A.; Motiwalla, Luvai; and Jonker, Catholijn M., "Microlearning mApp to Improve Long Term Health Behaviours: Design and Test of Multi-Channel Service Mix" (2014). *BLED 2014 Proceedings*. 4. http://aisel.aisnet.org/bled2014/4

This material is brought to you by the BLED Proceedings at AIS Electronic Library (AISeL). It has been accepted for inclusion in BLED 2014 Proceedings by an authorized administrator of AIS Electronic Library (AISeL). For more information, please contact elibrary@aisnet.org.

27th Bled eConference

eEcosystems

June 1 - 5, 2014; Bled, Slovenia

Microlearning mApp to Improve Long Term Health Behaviours: Design and Test of Multi-Channel Service Mix

Luuk P.A. Simons

Delft University of Technology, Netherlands
L.P.A.Simons@tudelft.nl

Florian Foerster

Georgia Institute of Technology, USA florian.foerster@gatech.edu

Peter A. Bruck

Research Studio Micro-learning and Information Environment; Research Studios

Austria Forschungsgesellschaft, Austria

bruck@researchstudio.at

Luvai Motiwalla

Operations & Info Systems Dept, U Massachusetts Lowell, USA luvai motiwalla@uml.edu

Catholijn M. Jonker

Delft University of Technology, Netherlands
C.M.Jonker@tudelft.nl

Abstract

Work place health support interventions are relevant: to support our aging work force and to help reduce the private and social burdens of preventable disease. Moreover, healthy living has become a priority in Dutch society.

Previous research showed that health apps should offer enough newness and relevance each time they are used. Otherwise the 'eHealth law of attrition' applies: 90% of users are lost prematurely.

We developed a hybrid health support solution that uses a mix of electronic and physical support services for improving health behaviours, including a mobile microlearning health quiz. This solution was tested in a multiple-case study at three work sites. The research question was whether a mobile health quiz can provide added value for users within a hybrid service mix and whether it promotes long term health readiness (awareness, motivation, plans and behaviours)?

We found that both the mobile health quiz and the overall hybrid solution contributed to health readiness and health behaviour improvements. Users indicated that the microlearning health quiz courses provided new and relevant information. We also observed relatively high utilization rates of the health quiz. Participants indicated that health behaviour insights were given that were directly applicable and that fitted well within the overall service mix. Moreover, even after 10 months they indicated to still have increased health awareness, -motivation and -behaviours (food, physical activity, mental recuperation).

A design analysis is conducted regarding service mix efficacy; the mobile microlearning health quiz helped fulfil key requirements for designing ICT-enabled lifestyle interventions, largely in the way it was anticipated.

Keywords: mobile Health, mApp, service design, multi-channel services, healthy lifestyle intervention

1 Introduction

Work place health support interventions are very relevant: to support our aging work force and to help reduce the private and social burdens of preventable disease. There are two design challenges there: increasingly more support is required, and this has to be delivered at lower costs. Hence, there is an urgency to develop automated eHealth solutions to improve user's health.

Previous research highlighted the following requirements in the work context: solutions need to be efficient; they should limit their intrusion on the other work tasks (Bruck, Motiwalla, Foerster 2012); they should offer enough newness and relevance (Simons, Guldemond, Hampe 2013) each time the mobile health application is used. Otherwise the eHealth law of attrition applies: 90% of users are lost prematurely (Eysenbach 2005).

This eHealth law of attrition is an important design challenge. On the one hand, we know that many health plans of people fail and that with extra support, success rates improve. On the other hand, automated eHealth support is often the economically preferred way (especially in the future, given our aging population). But if eHealth applications generally lose 90% of their users within several usage instances, we are not adding enough value with eHealth.

In previous research (Simons, Hampe, Guldemond 2012) we showed that various mobile health applications, even when they are popular mApps from the App stores, still face challenges in delivering added value to existing health coach relationships. In

several of our real world user tests, the 'eHealth law of attrition' applied (Eysenbach 2005). In other words, a large majority of users (>90%) stopped using the mApps within two weeks and judged that they added little new information or value after a few initial usage events. Very similar patterns were found for a food choices app (CalorieTeller), physical activity (Runkeeper) and a stress management app (Pranayama). Two main design lesson conclusions were (Simons, Hampe, Guldemond 2013, 2014): First, part of the attractiveness of an mApp appears to be the 'newness factor': do I learn or experience something new? Second, it helps when the mApp is a core element of a larger service mix within an existing health coach relationship, see also Jimison 2008.

Moreover, as part of qualitative user feedback, we found a continuous hunger from health coach participants for new health information. It was decided to create a health coach service mix, where the processes of health education would be supported with a mobile health education system in the form of a micro-learning Health Quiz on multiple health behaviour themes (food, physical activity, mental energy, long term sustainable health behaviours).

The overall research question is: Can a mobile health quiz provide added value within a hybrid (electronic and face to face) health coach service mix and does it promote long term health readiness (awareness, motivation, plans and behaviours)? We conducted a multiple-case study in three different work site environments (employers) and addressed the research question via four sub-questions: First, does the mobile health quiz provide added value (usefulness, fun, positive triggers) with low barriers (ease of use, limited time burdens)? Second, can the mobile health quiz be integrated in effective health coach processes and overall service mix? Third, do the mobile health quiz and the overall service mix improve health readiness (awareness, motivation, plans and behaviour experiments)? Fourth, do the mobile health quiz and the overall service mix improve health behaviours, short term and long term?

2 Theory

For health coach solutions to be effective, they need to help improve health readiness as indicated in the HAPA (Health Action Process Approach) model (Schwarzer 2008, Lippke 2009) and i-change models (De Vries and Mudde 1998). Three important phases are distinguished. Barriers or motivators for change can reside in each of these phases, which are: awareness, motivation/intention, and practice (including planning, experiencing, coping, improving). And as an underlying theme self-efficacy is important in these models: can we support people in developing skills to live more healthily and with tactics to deal with challenges?

If we look at the design challenge of persuasive technology (Fogg 2002, 2009) for health, it was theorized and tested elsewhere that this challenge is not just located in the ICT design, but also in the design of the overall service scape, including health effects and coach relationship (Simons, Hampe, Guldemond 2014). It should generate positive, mutually reinforcing service experiences across communication channels and activate long term health motivation and -behaviours, in order to deliver long term results. This is reflected in the following design evaluation framework for health improvement ICT solutions (Simons, Hampe, Guldemond 2014), see Figure 1. It evaluates the impact of

the ICT-enabled intervention on health effectiveness, coaching performance and ICT value adding.




Figure 1: Basic requirements when designing ICT-supported healthy lifestyle interventions

To increase solution impact, a hybrid or multi-channel service mix is recommended (Sperling, Simons and Bouwman 2009, Simons, Steinfield and Bouwman 2002, Simons 2006, Simons and Hampe 2010, 2010b), combining electronic and face-to-face interactions. For example, face to face 'on site' coaching had as benefits; a richer service experience with the coach, with other participants and with a health focused 'service scape'; group support experiences (obtaining additional social support and cocreating service experiences together); learning from each other; health experiences in healthy food-, sports- and relaxation exercises. Disadvantages are: more (travel) time needed; less flexibility regarding when and where; and not everyone likes group sessions (Demark-Wahnefried 2007). Electronic and (semi-)automated coaching has as benefits: more time-efficient; more flexibility in when and where to have contact; very explicit monitoring of your own progress online; having status reports including 'next steps' commitments always online. Disadvantages are: the sensory-, emotional- and group experiences are more limited. Also, the 'service scape' in which people are immersed is only virtual, not physical. In summary, often a hybrid service mix has most to offer.

In such a hybrid service mix, mobile micro-learning could potentially offer a number of advantages: it uses a personal device that is available any time any place, it is efficient and can use idle time that is otherwise lost, and it is suited for just in time learning (Bruck, Motiwalla, Foerster 2012).

3 Methods and Materials

Regarding our design research approach, we follow the design cycle of (Vaishnavi & Kuechler, 2004): from problem awareness and solution suggestion to development, evaluation and conclusion. After reporting our multiple-case study results in section 4, we will discuss design lessons in section 5.

3.1 Intervention: Hybrid lifestyle intervention with mHealth App

The mHealth Health Quiz App was used within a hybrid (face-to-face and digital) service mix. The healthy lifestyle support service mix consisted of:

- Digital health behaviour surveys 'BRAVO' with automated personal feedback (on physical activity, smoking, alcohol, food and energy/recuperation), at 0, 1, 3 and 10 months.
- A 3-hour start up workshop (with about 30 participants)
- Personal action plan drawn up at the start workshop
- Asking participants to use weekly buddy contacts
- The mHealth Health Quiz App (available for Android, iOS, blackberry and laptops)
- Supporting health education materials via email
- A 2-hour repeat workshop after 1 month for answering questions and for (peer) education
- Twelve times a weekly health tip email to help maintain awareness and motivation

The Health Quiz App was based on the micro-learning principles and technology platform as outlined previously (Bruck, Motiwalla, Foerster 2012). Hence, education is conducted via brief learning cards, each containing a question, (mostly) multiple choice answer options, plus a brief explanation after each answering attempt. The learning cards are organized in a number of courses and participants can see how much they progressed within a course. The course content was designed in the following way. It consists of seven courses of about 20 questions each. The first three courses educate participants on the 'what' or basic knowledge of healthy food-, exercise- and mental energy behaviours. The fourth course provides education and examples on what determines long term health behaviour success. Courses five through seven educate participants on the 'how' or daily tactics of healthy food-, exercise- and mental energy behaviours.

Within the course content design, questions are ranked in a certain order to address consecutive learning objectives. First, basic knowledge and awareness are increased, then motivation and plan making, next there is support for daily activities and coping strategies, plus seeking new self-norms and self-identity. Participants are free to switch between courses in order to support 'just in time' learning based on their needs. The Health Quiz Apps were downloaded and activated for all users during the 3-hour start up workshop.

3.2 Multiple-Case Study in three employer organizations

From Feb to Dec 2013, a multiple-case study was conducted with three employers to evaluate real world impacts of the healthy lifestyle intervention in Dutch work settings. The employers were: a Municipality (n=30 participants), an Advocacy organization for Dutch senior citizens (n=26 participants, half of them volunteer workers) and a Care Provider (n=30 participants).

From the perspective of these three organizations, an important question was: does this lifestyle intervention generate health behaviour improvements in the short and longer

term? This was assessed with a standardized Dutch online 'BRAVO' survey (on physical activity, smoking, alcohol, food and energy/recuperation), at 0, 1, 3 and 10 months. Moreover, at months 1, 3 and 10 several additional health behaviour and health readiness (according to the stages of awareness, willingness, plans and actions) questions were asked on top of the brief BRAVO question set.

Next, a design evaluation survey was added at month 1, and the findings from this survey were discussed in the 1-month group workshops at each organization, to obtain qualitative insights into the why and how of the survey data and user experiences. The design evaluation survey addressed: 1) an evaluation of health promotion value of the different elements in the service mix, including the Health Quiz App, 2) the microlearning mApp usefulness and ease of use. Besides, logging was conducted of use and progression in the micro-learning mApp environment.

4 Results

The overall patterns were rather similar across the three employer organizations regarding: adoption, evaluation and effects of the healthy lifestyle intervention. Hence, we first describe the generic outcomes. After that, we report cross-case differences.

At the start, there were 86 participants (n=30/26/30 for Municipality, Advocacy and Care Provider respectively). The survey response rates were 74% (n=64) at 1 month, then 55% (n=47) at 3 months and 56% at 10 months (n=48 for the BRAVO questions, n=47 for the rest).

4.1 Generic outcomes across the three employer organizations

From the perspective of the three employer organizations, the main result is that health behaviours improve, as measured with the standardized Dutch BRAVO survey (on physical activity, smoking, alcohol, food and energy/recuperation). As expected, the main improvement in health behaviour patterns occurs in the first month. After that, the new pattern is relatively stable, at least for 10 months period we measured. This pattern was observed within each organization. Table 1 provides the behaviour distribution accumulated for the three organizations at the start and 10 months. Table 2 summarizes the number of worsened and improved scores at months 1 and 10, compared to the start. Green means: compliant with Dutch health norms, yellow is nearly compliant and red is the rest.

Health Behaviours:	BRAVO survey results at start (n=86)	Percentages
Moderate Physical Activity		(40% - 9% - 51%)
Intensive Physical Activity		(22% - 26% - 52%)
Smoking		(83% - 0% - 17%)
Alcohol		(64% - 0% - 36%)
Food, Vegetables		(21% - 21% - 58%)




Table 1: Health behaviours distribution at start and 10 months. Green = health norm compliant. Yellow = nearly compliant. Red = the rest.

	Number of people, at 1 month (n=64)		Number of people, at 10 months (n=48)	
Health Behaviours:	Worsened:	Improved:	Worsened:	Improved:
Moderate Physical Activity	4	19	4	14
Intensive Physical Activity	4	21	4	18
Smoking	2	2	1	3
Alcohol	4	8	1	8
Food, Vegetables	2	23	3	16
Food, Fruits	5	32	1	18
Food, BMI:	2	4	1	5
Recuperation/Relaxing	5	16	3	14
Energy	5	17	5	14

Table 2: Summary of worsened and improved behaviour scores at 1 and 10 months (irt start)

Additional behaviour improvements reported in the surveys were (we list how many of the n=47 respondents (strongly) agreed at 10 months): I am physically active more often during the day (72%), I take a relax moment more often (47%), I more structurally

reduce my stress sources (51%), I eat fewer sugars and refined carbohydrates (55%), I eat more whole meal plant foods like nuts, mushrooms, olives etc (81%), I eat less red and processed meat (57%), I eat fewer butter fats (60%). Next, the health readiness indicators improved (awareness, intentions and plans), plus health effects in terms of improved physical or mental fitness. The scores at 1, 3 and 10 months were very similar. Table 3 provides the 10 months results (n=47).




Table 3: Health readiness and fitness improvements after 10 months, n=47. Green = (highly) agree. Yellow = neutral. Red/orange = (highly) disagree.

Next, a question was: how did the Health Quiz App and other service mix elements contribute to the results above? This question was part of the 1-month evaluation. In the survey data as well as the workshops observed that not everybody had preferences for the same service elements. Still, most service elements were judged to be helpful for improving health behaviours by a significant part of the users, see Table 4. The lowest helpfulness scores as percentage '(strongly) agree', n=64, were given for: doing it as a group (20%), having a buddy (25%) and the weekly health tip mail (36%).

The highest helpfulness scores as percentage '(strongly) agree', n=64, were given for: having a second workshop at 1 month (52%), feeling better (53%), the Health Quiz mApp (53%), making my own health activity plan (66%), knowing what my own influence is (75%) and the start workshop (77%).






Table 4: Service elements that stimulated healthier behaviours (n=64, at 1 month). Green = (highly) agree. Yellow = neutral. Red/orange = (highly) disagree. Grey = not applicable.

A next set of evaluation questions at 1 month regarded usefulness, added value and ease of use of the micro-learning Health Quiz mApp. The majority of respondents indicated that it was efficient, useful, fun, that they would have learned less without it and that they now made healthier choices thanks to the micro-learning Health Quiz mApp. Likewise, the majority also indicated that it fulfilled a desire to learn more after the start workshop, that it was low effort, that the (smaller) mobile screens were no hindrance, that most learning cards were relevant, that things were learned that were directly applicable and that the courses provided regular stimuli to make healthy choices. These findings were confirmed in the feedback during the 1-month workshops.

Moreover, the qualitative findings were corroborated via the micro-learning logging data, in the sense that the majority of participants who started the micro-learning Health Quiz mApp also completed all 7 available courses. In contrast with the eHealth law of attrition (Eysenbach 2005), completion rates were relatively high. On an intention to treat basis, 66% of the courses were completed that were made available for the n=86 starting participants. In the Municipality case (n=30), 14 participants completed all 7 courses, five completed several courses and 11 completed none (of which 7 participants never started the micro-training). The total completion rate was 53% of all available course material. In the Advocacy case (n=26), 18 participants completed all 7 courses, and 8 completed none (of which 5 never started the micro-training). Their total completion rate was 69% of all available course material. In the Care Provider case (n=30), 21 participants completed all 7 courses, four completed several courses and five completed none (all of them did start the micro-training and completed multiple questions, mostly sampling them across multiple courses). The total completion rate was 75% of all available course material.

4.2 Cross-case differences

There were a few case-specific characteristics that had an impact on adoption patterns and compliance and response rates. However, the overall design evaluation findings were very similar across cases. In Table 3 we highlight the differences.

Employer case	Case-specific characteristics and findings	
Municipality	 Relatively lower start- and completion rates, partly due to the fact that several participants were sent by their managers and did not participate on a voluntary basis. For those who did participate, the relative improvements in eating vegetables and adding moderate intensity physical activity was larger than in the other cases. 	
Advocacy	 About half of the group were (senior) volunteers and the average age in this group was highest. Several sudden events happened in the lives of these participants, hampering workshop participation and course completion for several of them. Initial health behaviour scores were highest in this group: for daily physical 	

	activity, for fruit and vegetables consumption. However, their energy was lower and stayed low. - About 50% were retired: some were more busy than ever. Others reported that the topics of work related stress and energy were less relevant for them.
Care Provider	 The regional director was a strong health advocate, participant and initiator in this group. Start- and compliance rates were highest is this group. This group had the lowest fruit consumption start, but the highest final score. The improvements were largest in energy and recuperation behaviours were largest in this group, as well as the reported gain in mental fitness (at 1, 3 and 10 months).

Table 5: Cross-case differences

5 Discussion and Conclusion

In conclusion, the short answer to all four research questions in this paper is yes, and we will discuss them in relation to the evaluation framework of Figure 1. First, the mobile health quiz provides added value (usefulness, fun, positive triggers) with low barriers (ease of use, limited time burdens) Second, according to the participants the mobile health quiz is well integrated in the overall service mix. Hence, the 'ICT effectiveness' factor of Figure 1 appears to be sufficiently addressed. The mobile health quiz also increases the 'coaching effectiveness' factor of Figure 1 by providing regular triggers for health awareness, coping strategies and useful health behaviours. In answer to the third research sub-question, health readiness is improved: awareness, motivation, plans and behaviour experiments. Fourth, the various health behaviours are improved from the Dutch BRAVO survey (on physical activity, smoking, alcohol, food and energy/recuperation).

The answers to the third and fourth research sub-questions demonstrate the contribution to the 'health effectiveness' factor of Figure 1, both of the overall hybrid service mix and of the mobile micro-learning health quiz within that mix. Besides explicit participant feedback, we have the indications from the course completion rates of the mobile health quiz, which were 66%: well above the Eysenbach (2005) 'law'. Given the fact that these courses were not mandatory for these participants, but only additional support for health self-management, and given the fact that each course easily takes 20 minutes to complete (in a context of time scarcity, Bruck, Motiwalla, Foerster 2012) we regard course completion rates as a sensible proxy for perceived usefulness.

Health behaviours not only improved at 1 and 3 months, but also 10 months after the start. The latter finding is relatively special, in the sense that most healthy lifestyle interventions, whether at work sites (Verweij, Coffeng, Mechelen, Proper 2011) or not (Seidell and Halberstadt 2011), tend to generate only short term results (3 or 6 months), after which people generally fall back into their old patterns.

Our main *contribution to theory* is twofold. In this empirical test of our proof of concept with three organisations we see a tentative confirmation of the two propositions on which we have built our design research. First, a mobile micro-learning health quiz appears useful for fulfilling the three key design requirements when designing ICT-supported healthy lifestyle interventions: health-, coaching- and ICT effectiveness. And

second, as contributions on these three requirements increased, we indeed empirically observed an improved health readiness. Thus, our proposition was confirmed that the requirements from Figure 1 are important for ICT-enabled health intervention success.

Regarding *practical implications*, the relevant design question is what made these results come about and how can we improve even further? In this section we first discuss study limitations and then conduct a design evaluation, using the framework from our theory section.

This study has several limitations. First, it is largely qualitative, evaluating effects across three case organizations. In the survey, users mostly agree on issues of microlearning usefulness and ease of use. Statistical techniques like explorative regression analysis were not possible, since the variance in answers was too low. Second, our survey results are likely subject to self-selection biases: our survey respondents are largely those users who completed most or all of the micro-learning courses. They are the ones most likely to be positive. Third, the context is different in each case organization and in our study design we cannot control for confounding factors. On the other hand, the strong evaluation- and effect agreement between participants across organizations does hint at the robustness and cross-case validity of the findings. For a more rigorous evaluation, we are preparing an experimental design to compare intervention and control groups, where the latter receive the service mix without the mobile micro-learning health quiz.

Table 6 shows that in our own authors' opinions there are several points for improvement. Regarding factor 1 of Figure 1, health effectiveness, more objective improvement measures could be advantageous. Especially for self-management of people with health issues (e.g. diabetes-2, irregular high blood pressure or heart arrhythmia, or impaired renal function) 24x7 monitoring of effects of lifestyle improvement can be very beneficial. For the second factor, coaching effectiveness, there is a challenge of automatically integrating context- and health information. For example, if we notice that someone has been sitting most of the day, when will reminders/triggers be appreciated to get up and move about, and when not? For example, if I'm very busy with finishing a report or having urgent meetings, it can be a conscious and preferred strategy to continue a sedentary work activity for the time being. Reminders and triggers can easily become a nuisance. But at other times, when falling in my coach potato trap in the evening, I may very well appreciate more persistent triggers to go play sports with a buddy. Finally, ICT value adding (factor 3) could be improved via at least two routes. First, improving automated logging of health (behaviour) data and integrating this into coach processes. Second, designing more intelligent, interactive coach processes, which incorporate user preferences and plans, contextual/situational priorities and health data consequences.

Health Effectiveness	Coaching Performance	ICT Value Adding
++ Health Literacy: Impacts	+/- Promoting health actions:	+ Motivators, triggers,
from Health Quiz, workshops	Many health tips are provided.	experiences: health quiz, mail
and education materials.	Impact depends on execution	tips and surveys provided
+ Health behaviours: BRAVO	of plans.	triggers, (fun) experiences plus
survey indicates improvements.	++ Supporting self-efficacy:	hope and improvement

+/- Health outcomes: Feeling	Users indicate a strong	opportunities.
more fit is a positive outcome,	contribution from 'know what	+/- Simplicity: Installation and
but more objective measures	my own influence is'	first use were burdening for
not used.	++ Activating intrinsic	some. After that usage was
+ Quality of Life: Feeling	motivation: A strongly activated	simple, low effort.
better, mentally and physically	desire to improve, plus rewards	+/- Fit with coach processes:
more fit.	via feeling better.	Users felt synergies with the
		workshops, education
		materials, personal action plans
		and answering individual
		questions.
Potential improvement:	Potential improvement:	Potential improvement:
Using more objective health	Context aware and	Coach processes could be
outcomes, possibly future 24x7	personalized coaching.	automated more (e.g.
health tracking.		goals/means support).

Table 6: Design evaluation (authors' opinions, 5-point scale from - - to ++)

In summary, given the relatively static content of the micro-learning health quiz, it served its health support goals well, thanks to the other service mix elements and the overall service concept. eCoach improvement opportunities for the future abound, of which we identified several.

References

- Baumeister RF, Tierney J. (2011). Willpower: Rediscovering the Greatest Human Strength. New York: Penguin Group.
- Bruck PA, Motiwalla L, Foerster F. 2012. Mobile Learning with Micro-content: A Framework and Evaluation. Paper presented at the 25th Bled eConference. from www.bledconference.org.
- Daubenmier, J., Weidner, G., Marlin, R., Crutchfield, L. and S. Dunn-Emke, C. C., B. Gao, P. Carroll, D. Ornish. (2006). Lifestyle and health-related quality of life of men with prostate cancer managed with active surveillance. Urology, 67(1), pp. 125-130.
- Demark-Wahnefried, W., Clipp, E., Lipkus, I., Lobach, D. et al. (2007). Main Outcomes of the FRESH START Trial: A Sequentially Tailored, Diet and Exercise Mailed Print Intervention Among Breast and Prostate Cancer Survivors. J Clin Oncol, 25(19), pp. 2709-2718.
- De Vries, H. and Mudde, A. (1998). Predicting stage transitions for smoking cessation applying the Attitude Social influence Efficacy Model. Psychology & Health, 13, 369–385
- Eysenbach G. 2005. The Law of Attrition. J Med Internet Res. 7(1): e11.
- Fogg, B.J. (2002). Persuasive technology: using computers to change what we think and do." Ubiquity, December (2002): 5.

- Fogg, B. J. (2009). A behavior model for persuasive design. Proceedings of the 4th international conference on persuasive technology. ACM, 2009.
- Jimison, H., Gorman, P., Woods, S., Nygren, P., Walker, M., et al. (2008). Barriers and drivers of health information technology use for the elderly, chronically ill, and underserved. Evid Rep Technol Assess (Full Rep) Nov;(175):1–1422.
- Lippke, S., Wiedemann, A. U., Ziegelmann, J. P., Reuter, T. and Schwarzer, R. (2009). Self-efficacy moderates the mediation of intentions into behavior via plans. American Journal of Health Behavior, 33(5), 521–529.
- Ornish, D. (2008). The Spectrum: A Scientifically Proven Program to Feel Better, Live Longer, Lose Weight, Gain Health. New York: Ballantine.
- Schwarzer, R. (2008). Modeling health behavior change: How to predict and modify the adoption and maintenance of health behaviors. Applied Psychology: An International Review, 57(1), 1–29.
- Seidell J, Halberstadt J (2011) Tegenwicht: Feiten en Fabels over overgewicht, Bert Bakker, Amsterdam, pp232.
- Simons, LPA and Bouwman, H. (2004). Designing a click and mortar channel mix. *International Journal of Internet Marketing and Advertising* 1(3): 229–250.
- Simons, LPA. (2006). *Multi-channel services for click and mortars: development of a design method*. PhD Thesis, Delft University of Technology.
- Simons, LPA, and Hampe, JF. (2010). Service Experience Design for Healthy Living Support; Comparing an In-House with an eHealth Solution. Paper *presented at the 23rd Bled eConference*. Bled, Slovenia, from www.bledconference.org.
- Simons, LPA and Hampe, JF. (2010b). Exploring e/mHealth Potential for Health Improvement; A Design Analysis for Future e/mHealth Impact. Paper *presented* at the 23rd Bled eConference. Bled, Slovenia, from www.bledconference.org.
- Simons, LPA, Hampe JF, and Guldemond NA. (2012). Designing Healthy Consumption Support: Mobile application use added to (e)Coach Solution. Paper *presented at the 25th Bled eConference*. Bled, Slovenia, from www.bledconference.org.
- Simons LPA, Hampe JF, Guldemond NA. (2013). Designing Healthy Living Support: Mobile applications added to hybrid (e)Coach Solution, *Health and Technology*, 3 (1), pp.1-11.
- Simons LPA, Hampe JF, Guldemond NA. (2014). Designing ICT-support for healthy lifestyle interventions. To appear in *Electronic Markets*. DOI 10.1007/s12525-014-0157-7
- Simons, LPA, Steinfield C and Bouwman H. (2002) "Strategic positioning of the Web in a multi-channel market approach." *Internet Research* 12 (4): 339-347.
- Simons, LPA and Verhagen WP. (2008). Applying value-sensitive design and quality function deployment to healthcare ICT: the case of Dutch primary care unit dossiers. Journal of Design Research 7 (2): 155-176.

- Sperling, R., L.P.A. Simons and H. Bouwman. (2009). Multi-Channel Service Concept Definition and Prototyping, *International Journal of Electronic Business*, 7 (3), pp.237–255.
- Vaishnavi, V and Kuechler, W. 2004. Design Research in Information Systems. Last updated August 16, 2009 from http://desrist.org/design-research-in-information-systems
- Verweij, LM, Coffeng, J., van Mechelen, W., Proper, KI (2011). Meta-analyses of workplace physical activity and dietary behaviour interventions on weight outcomes. Obesity Reviews, 12, 406-429.